

This example is set up in ECF Augie (with CM symbols and Euler Greek). It uses:

```
\renewcommand\familydefault{fau} % emerald package  
\usepackage[defaultmathsizes,eulergreek]{mathastext}
```

Typeset with mathastext l.15c (2012/10/05).

To illustrate some Hilbert Space properties of the co-Poisson summation, we will assume $K = \mathbb{Q}$. The components (a_v) of an adèle a are written a_p at finite places and a_r at the real place. We have an embedding of the Schwartz space of test-functions on \mathbb{R} into the Bruhat-Schwartz space on A which sends $\psi(x)$ to $\varphi(a) = \prod_p |a_p|_p \leq 1 (a_p) \cdot \psi(a_r)$, and we write $E'_R(G)$ for the distribution on \mathbb{R} thus obtained from $E'(G)$ on A .

Theorem 1. Let G be a compact Bruhat-Schwartz function on the ideles of \mathbb{Q} . The co-Poisson summation $E'_R(G)$ is a square-integrable function (with respect to the Lebesgue measure). The $L^2(\mathbb{R})$ function $E'_R(G)$ is equal to the constant $-\int_{A^\times} G(v) |v|^{-1/2} d^*v$ in a neighborhood of the origin.

Proof. We may first, without changing anything to $E'_R(G)$, replace G with its average under the action of the finite unit ideles, so that it may be assumed invariant. Any such compact invariant G is a finite linear combination of suitable multiplicative translates of functions of the type $G(v) = \prod_p |v_p|_p = (v_p) \cdot f(v_r)$ with $f(t)$ a smooth compactly supported function on \mathbb{R}^\times , so that we may assume that G has this form. We claim that:

$$\int_{A^\times} |\varphi(v)| \sum_{Q \in \mathbb{Q}^\times} |G(Qv)| \sqrt{|v|} d^*v < \infty$$

Indeed $\sum_{Q \in \mathbb{Q}^\times} |G(Qv)| = |f(|v|)| + |f(-|v|)|$ is bounded above by a multiple of $|v|$. And $\int_{A^\times} |\varphi(v)| |v|^{3/2} d^*v < \infty$ for each Bruhat-Schwartz function on the adeles (Basically, from $\prod_p (1-p^{-3/2})^{-1} < \infty$). So

$$E'(G)(\varphi) = \sum_{Q \in \mathbb{Q}^\times} \int_{A^\times} \varphi(v) G(Qv) \sqrt{|v|} d^*v - \int_{A^\times} \frac{G(v)}{\sqrt{|v|}} d^*v \int_A \varphi(x) dx$$

$$E'(G)(\varphi) = \sum_{Q \in \mathbb{Q}^\times} \int_{A^\times} \varphi(v/Q) G(v) \sqrt{|v|} d^*v - \int_{A^\times} \frac{G(v)}{\sqrt{|v|}} d^*v \int_A \varphi(x) dx$$

Let us now specialize to $\varphi(a) = \prod_p |a_p|_p \leq 1 (a_p) \cdot \psi(a_r)$. Each integral can be evaluated as an infinite product. The finite places contribute 0 or 1 according to whether $Q \in \mathbb{Q}^\times$

satisfies $\|q\|_p < 1$ or not. So only the inverse integers $q = 1/n, n \in \mathbb{Z}$, contribute:

$$E'_R(G)(\psi) = \sum_{n \in \mathbb{Z}^\times} \int_{\mathbb{R}^\times} \psi(nt) f(t) \sqrt{|t|} \frac{dt}{2|t|} - \int_{\mathbb{R}^\times} \frac{f(t)}{\sqrt{|t|}} \frac{dt}{2|t|} \int_{\mathbb{R}} \psi(x) dx$$

We can now revert the steps, But this time on \mathbb{R}^\times and we get:

$$E'_R(G)(\psi) = \int_{\mathbb{R}^\times} \psi(t) \sum_{n \in \mathbb{Z}^\times} \frac{f(t/n)}{\sqrt{|n|}} \frac{dt}{2\sqrt{|t|}} - \int_{\mathbb{R}^\times} \frac{f(t)}{\sqrt{|t|}} \frac{dt}{2|t|} \int_{\mathbb{R}} \psi(x) dx$$

Let us express this in terms of $\alpha(y) = (f(y) + f(-y))/2\sqrt{|y|}$:

$$E'_R(G)(\psi) = \int_{\mathbb{R}} \psi(y) \sum_{n \geq 1} \frac{\alpha(y/n)}{n} dy - \int_0^\infty \frac{\alpha(y)}{y} dy \int_{\mathbb{R}} \psi(x) dx$$

So the distribution $E'_R(G)$ is in fact the even smooth function

$$E'_R(G)(y) = \sum_{n \geq 1} \frac{\alpha(y/n)}{n} - \int_0^\infty \frac{\alpha(y)}{y} dy$$

As $\alpha(y)$ has compact support in $\mathbb{R} \setminus \{0\}$, the summation over $n \geq 1$ contains only vanishing terms for $|y|$ small enough. So $E'_R(G)$ is equal to the constant $-\int_0^\infty \frac{\alpha(y)}{y} dy = -\int_{\mathbb{R}^\times} \frac{f(y)}{\sqrt{|y|}} \frac{dy}{2|y|} = -\int_{\mathbb{A}^\times} g(t)/\sqrt{|t|} d^*t$ in a neighborhood of 0. To prove that it is L^2 , let $\beta(y)$ be the smooth compactly supported function $\alpha(1/y)/2|y|$ of $y \in \mathbb{R}$ ($\beta(0) = 0$). Then ($y \neq 0$):

$$E'_R(G)(y) = \sum_{n \in \mathbb{Z}} \frac{1}{|y|} \beta\left(\frac{n}{y}\right) - \int_{\mathbb{R}} \beta(y) dy$$

From the usual Poisson summation formula, this is also:

$$\sum_{n \in \mathbb{Z}} \gamma(ny) - \int_{\mathbb{R}} \beta(y) dy = \sum_{n \neq 0} \gamma(ny)$$

where $\gamma(y) = \int_{\mathbb{R}} \exp(i2\pi yw) \beta(w) dw$ is a Schwartz rapidly decreasing function. From this formula we deduce easily that $E'_R(G)(y)$ is itself in the Schwartz class of rapidly decreasing functions, and in particular it is square-integrable. \square

It is useful to recapitulate some of the results arising in this proof:

Theorem 2. Let g be a compact Bruhat-Schwartz function on the ideles of \mathbb{Q} . The co-Poisson summation $E'_R(g)$ is an even function on \mathbb{R} in the Schwartz class of rapidly decreasing functions. It is constant, as well as its Fourier Transform, in a neighborhood of the origin. It may be written as

$$E'_R(g)(y) = \sum_{n \neq 0} \frac{\alpha(y/n)}{n} - \int_0^\infty \frac{\alpha(y)}{y} dy$$

with a function $\alpha(y)$ smooth with compact support away from the origin, and conversely each such formula corresponds to the co-Poisson summation $E'_R(g)$ of a compact Bruhat-Schwartz function on the ideles of \mathbb{Q} . The Fourier transform $\int_{\mathbb{R}} E'_R(g)(y) \exp(i2\pi wy) dy$ corresponds in the formula above to the replacement $\alpha(y) \mapsto \alpha(1/y)/|y|$.

Everything has been obtained previously.