

This example is set up in GNU FreeFont Serif with TX fonts symbols and delimiters.

```
\usepackage[no-math]{fontspec}
\usepackage{txfonts} %\let\mathbb=\varmathbb
\setmainfont[ExternalLocation,
    Mapping=tex-text,
    BoldFont=FreeSerifBold,
    ItalicFont=FreeSerifItalic,
    BoldItalicFont=FreeSerifBoldItalic]{FreeSerif}
\usepackage[defaultmathsizes]{mathastext}
```

Typeset with mathastext 1.15d (2012/10/13).
(compiled with X_EL^AT_EX)

Theorem 1. Let there be given indeterminates $u_i, v_i, k_i, x_i, y_i, l_i$, for $1 \leq i \leq n$. We define the following $n \times n$ matrices

$$U_n = \begin{pmatrix} u_1 & u_2 & \dots & u_n \\ k_1 v_1 & k_2 v_2 & \dots & k_n v_n \\ k_1^2 u_1 & k_2^2 u_2 & \dots & k_n^2 u_n \\ \vdots & \dots & \dots & \vdots \end{pmatrix} \quad V_n = \begin{pmatrix} v_1 & v_2 & \dots & v_n \\ k_1 u_1 & k_2 u_2 & \dots & k_n u_n \\ k_1^2 v_1 & k_2^2 v_2 & \dots & k_n^2 v_n \\ \vdots & \dots & \dots & \vdots \end{pmatrix} \quad (1)$$

where the rows contain alternatively u 's and v 's. Similarly:

$$X_n = \begin{pmatrix} x_1 & x_2 & \dots & x_n \\ l_1 y_1 & l_2 y_2 & \dots & l_n y_n \\ l_1^2 x_1 & l_2^2 x_2 & \dots & l_n^2 x_n \\ \vdots & \dots & \dots & \vdots \end{pmatrix} \quad Y_n = \begin{pmatrix} y_1 & y_2 & \dots & y_n \\ l_1 x_1 & l_2 x_2 & \dots & l_n x_n \\ l_1^2 y_1 & l_2^2 y_2 & \dots & l_n^2 y_n \\ \vdots & \dots & \dots & \vdots \end{pmatrix} \quad (2)$$

There holds

$$\det_{1 \leq i, j \leq n} \left(\frac{u_i y_j - v_i x_j}{l_j - k_i} \right) = \frac{1}{\prod_{i,j} (l_j - k_i)} \begin{vmatrix} U_n & X_n \\ V_n & Y_n \end{vmatrix}_{2n \times 2n} \quad (3)$$

Proof. Let A, B, C, D be $n \times n$ matrices, with A and C invertible. Using $\begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} A & 0 \\ 0 & C \end{pmatrix} \begin{pmatrix} I & A^{-1}B \\ I & C^{-1}D \end{pmatrix}$ we obtain

$$\begin{vmatrix} A & B \\ C & D \end{vmatrix} = |A||C||C^{-1}D - A^{-1}B| \quad (4)$$

where vertical bars denote determinants. Let $d(u) = \text{diag}(u_1, \dots, u_n)$ and $p_u = \prod_{1 \leq i \leq n} u_i$. We define similarly $d(v), d(x), d(y)$ and p_v, p_x, p_y . From the previous identity we get

$$\begin{aligned} \begin{vmatrix} Ad(u) & Bd(x) \\ Cd(v) & Dd(y) \end{vmatrix} &= |A||C| p_u p_v \begin{vmatrix} d(v)^{-1} C^{-1} D d(y) - d(u)^{-1} A^{-1} B d(x) \end{vmatrix} \\ &= |A||C| \begin{vmatrix} d(u) C^{-1} D d(y) - d(v) A^{-1} B d(x) \end{vmatrix} \end{aligned} \quad (5)$$

The special case $A = C, B = D$, gives

$$\begin{vmatrix} Ad(u) & Bd(x) \\ Ad(v) & Bd(y) \end{vmatrix}_{2n \times 2n} = \det(A)^2 \det_{1 \leq i, j \leq n} ((u_i y_j - v_i x_j)(A^{-1}B)_{ij}) \quad (6)$$

Let $W(k)$ be the Vandermonde matrix with rows $(1 \dots 1), (k_1 \dots k_n), (k_1^2 \dots k_n^2), \dots$, and $\Delta(k) = \det W(k)$ its determinant. Let

$$K(t) = \prod_{1 \leq m \leq n} (t - k_m) \quad (7)$$

and let C be the $n \times n$ matrix $(c_{im})_{1 \leq i, m \leq n}$, where the c_{im} 's are defined by the partial fraction expansions:

$$1 \leq i \leq n \quad \frac{t^{i-1}}{K(t)} = \sum_{1 \leq m \leq n} \frac{c_{im}}{t - k_m} \quad (8)$$

We have the two matrix equations:

$$C = W(k) \operatorname{diag}(K'(k_1)^{-1}, \dots, K'(k_n)^{-1}) \quad (9a)$$

$$C \cdot \left(\frac{1}{l_j - k_m} \right)_{1 \leq m, j \leq n} = W(l) \operatorname{diag}(K(l_1)^{-1}, \dots, K(l_n)^{-1}) \quad (9b)$$

This gives the (well-known) identity:

$$\left(\frac{1}{l_j - k_m} \right)_{1 \leq m, j \leq n} = \operatorname{diag}(K'(k_1), \dots, K'(k_n)) W(k)^{-1} W(l) \operatorname{diag}(K(l_1)^{-1}, \dots, K(l_n)^{-1}) \quad (10)$$

We can thus rewrite the determinant we want to compute as:

$$\left| \frac{u_i y_j - v_i x_j}{l_j - k_i} \right|_{1 \leq i, j \leq n} = \prod_m K'(k_m) \prod_j K(l_j)^{-1} \left| (u_i y_j - v_i x_j) (W(k)^{-1} W(l))_{ij} \right|_{n \times n} \quad (11)$$

We shall now make use of (6) with $A = W(k)$ and $B = W(l)$.

$$\begin{aligned} \left| \frac{u_i y_j - v_i x_j}{l_j - k_i} \right|_{1 \leq i, j \leq n} &= \Delta(k)^{-2} \prod_m K'(k_m) \prod_j K(l_j)^{-1} \begin{vmatrix} W(k)d(u) & W(l)d(x) \\ W(k)d(v) & W(l)d(y) \end{vmatrix} \\ &= \frac{(-1)^{\frac{n(n-1)}{2}}}{\prod_{i,j} (l_j - k_i)} \begin{vmatrix} W(k)d(u) & W(l)d(x) \\ W(k)d(v) & W(l)d(y) \end{vmatrix}_{2n \times 2n} \end{aligned} \quad (12)$$

The sign $(-1)^{n(n-1)/2} = (-1)^{[\frac{n}{2}]}$ is the signature of the permutation which exchanges rows i and $n+i$ for $i = 2, 4, \dots, 2[\frac{n}{2}]$ and transforms the determinant on the right-hand side into $\begin{vmatrix} U_n & X_n \\ V_n & Y_n \end{vmatrix}$. This concludes the proof. \square